

# Were last glacial climate events simultaneous between Greenland and France? A quantitative comparison using non-tuned chronologies

MAARTEN BLAAUW,<sup>1\*</sup> BARBARA WOHLFARTH,<sup>2</sup> J. ANDRÉS CHRISTEN,<sup>3</sup> LINDA AMPEL,<sup>4</sup> DANIEL VERES,<sup>4,5</sup> KONRAD A. HUGHEN,<sup>6</sup> FRANK PREUSSER<sup>7</sup> and ANDERS SVENSSON<sup>8</sup>

<sup>1</sup> School of Geography, Archaeology and Palaeoecology, Queen's University Belfast, UK

<sup>2</sup> Department of Geology and Geochemistry, Stockholm University, Stockholm, Sweden

<sup>3</sup> Centro de Investigación en Matemáticas CIMAT, Guanajuato, Mexico

<sup>4</sup> Department of Physical Geography and Quaternary Geology, Stockholm University, Stockholm, Sweden

<sup>5</sup> 'Emil Racovita' Institute of Speleology, Cluj-Napoca, Romania

<sup>6</sup> Woods Hole Oceanographic Institution, Woods Hole, MA, USA

<sup>7</sup> Institut für Geologie, Universität Bern, Bern, Switzerland

<sup>8</sup> Ice and Climate Research, Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark

Blaauw, M., Wohlfarth, B., Christen, J. A., Ampel, L., Veres, D., Hughen, K. A., Preusser, F. and Svensson, A. 2010. Were last glacial climate events simultaneous between Greenland and France? A quantitative comparison using non-tuned chronologies. *J. Quaternary Sci.*, Vol. 25 pp. 387–394. ISSN 0267-8179.

Received 30 March 2009; Revised 4 June 2009; Accepted 5 June 2009

**ABSTRACT:** Several large abrupt climate fluctuations during the last glacial have been recorded in Greenland ice cores and archives from other regions. Often these Dansgaard–Oeschger events are assumed to have been synchronous over wide areas, and then used as tie-points to link chronologies between the proxy archives. However, it has not yet been tested independently whether or not these events were indeed synchronous over large areas. Here, we compare Dansgaard–Oeschger-type events in a well-dated record from southeastern France with those in Greenland ice cores. Instead of assuming simultaneous climate events between both archives, we keep their age models independent. Even these well-dated archives possess large chronological uncertainties that prevent us from inferring synchronous climate events at decadal to multi-centennial time scales. If possible, comparisons between proxy archives should be based on independent, non-tuned time-scales. Copyright © 2009 John Wiley & Sons, Ltd.



## Introduction

Some of the most intriguing climatic features of the last glacial period are the large abrupt temperature fluctuations that recurred at millennial timescales. These Dansgaard–Oeschger (D-O) events, originally seen in  $\delta^{18}\text{O}$  variations in Greenland ice cores (Dansgaard *et al.*, 1993), have subsequently been reported from proxy archives around the North Atlantic (Bond *et al.*, 1993; Allen *et al.*, 1999; Cacho *et al.*, 1999; Shackleton *et al.*, 2000; Sánchez-Goñi *et al.*, 2002; Voelker *et al.*, 2002; Tzedakis *et al.*, 2004; Roucoux *et al.*, 2005; Andersen *et al.*, 2006; Vautravers and Shackleton, 2006; Wohlfarth *et al.*, 2008) and beyond (Wang *et al.*, 2001; Hendy *et al.*, 2002; Voelker *et al.*, 2002; Burns *et al.*, 2003; Rohling *et al.*, 2003; Hughen *et al.*, 2004; Turney *et al.*, 2004). D-O events have been hypothesised to result from major shifts in North Atlantic meridional overturning circulation (Knutti *et al.*, 2004; EPICA

Community Members, 2006). A rapid atmospheric transmission of the signal could have led to more or less synchronous D-O style events within the Northern Hemisphere. Indeed, abrupt climatic events recognised in marine and terrestrial proxy time series are frequently reported to have been synchronous with Greenland D-O events (Bond *et al.*, 1993; Allen *et al.*, 1999; Cacho *et al.*, 1999; Wang *et al.*, 2001; Sánchez-Goñi *et al.*, 2002; Burns *et al.*, 2003; Hughen *et al.*, 2004; Roucoux *et al.*, 2005; Vautravers and Shackleton, 2006).

However, the inference of synchronicity between regions is constrained by numerous factors, including the response of the analysed marine or terrestrial proxies to local, regional and global climate signals, the temporal resolution of the studied sequence, and the applied age models. The lack of an internationally accepted radiocarbon ( $^{14}\text{C}$ ) calibration curve beyond 26 cal. ka BP (Reimer *et al.*, 2004; van der Plicht *et al.*, 2004), uncertain and time-dependent radiocarbon 'reservoir age' offsets for glacial marine records (Muschele *et al.*, 2008) and chronological uncertainties of up to 2600 a (at 2 standard deviations) in the Greenland ice cores (Andersen *et al.*, 2006) further add to this complexity. In order to overcome these difficulties, timescales and events in marine and terrestrial

\*Correspondence to: M. Blaauw, School of Geography, Archaeology and Palaeoecology, Queen's University, Belfast, UK.  
E-mail: maarten.blaauw@qub.ac.uk

records are often tuned (artificially synchronised) to ice core chronologies (e.g. Bond *et al.*, 1993; Cacho *et al.*, 1999; Walker *et al.*, 1999; Tzedakis *et al.*, 2004; Skinner, 2008).

Tuning assumes synchronicity of major climatic/environmental events between multiple proxy archives. According to the tuning line of reasoning, major climatic events were so large that they must have been expressed in a wide spatial range of fossil proxy records. These synchronous proxy changes are then used as tie-points (points of equal age) between the archives, thus providing age–depth estimates for one core based on the age model of a better dated ‘mother’ time series (e.g. orbital forcing, Greenland ice cores, Hulu cave speleothems or stacked marine  $\delta^{18}\text{O}$  curves; Bond *et al.*, 1993; Cacho *et al.*, 1999; Tzedakis *et al.*, 2004; Lisiecki and Raymo, 2005; Hughen *et al.*, 2006). A good example of a tie-point is the North Atlantic Ash Zone II tephra, resulting from a single volcanic eruption, and deposited over a wide area as it has been encountered and identified geochemically in Greenland ice as well as in a range of North Atlantic marine sediment cores (Austin *et al.*, 2004). This tephra, dated at ca.  $55\,330 \pm 1184$  cal. BP in NGRIP (Svensson *et al.*, 2008), was deposited during a major environmental event found both in marine and ice cores, and thus this event can safely be assumed to have been synchronous (interstadial 15, Bond *et al.*, 1993; Austin *et al.*, 2004). Even so, this tephra might have been produced by two volcanic eruptions separated by up to 2000 a (Wåstegård *et al.*, 2006).

Many other environmental events used to align glacial proxy archives are not accompanied by uniquely identifiable ‘fingerprints’. Instead, tie-points are often inferred based on the order, amplitudes and shapes (e.g. ‘tooth-like’ or ‘w-shaped’) of proxy curves, which in turn reflect the size and abruptness of climate change events (Bond *et al.*, 1993; Shackleton *et al.*, 2000; Tzedakis *et al.*, 2004). Besides the tephra tie-point discussed above, only few event-based glacial tie-points have been dated independently (e.g. using radiocarbon dates); dating densities were often low and discrepancies between tuning and independent dates were at millennial scales (Hendy *et al.*, 2002; Rohling *et al.*, 2003; Shackleton *et al.*, 2004; Svensson *et al.*, 2006; Skinner, 2008). The ‘mother’ chronologies themselves also possess considerable chronological uncertainties (e.g., 5000 a for SPECMAP; Martinson *et al.*, 1987), so any tuned ‘daughter’ chronology will necessarily obtain larger errors (a combination of the uncertainties of the mother chronology, of the tuning process, and of the uncertain sedimentation histories between tie-points). The assumption of synchronicity between regions eliminates any potential for objective evaluation of relative timing in different timescales (Walker *et al.*, 1999; Wunsch, 2006; Blaauw *et al.*, 2007a). Moreover, these climate-driven linkages are based on plausible yet unproven assumptions on regional to hemispheric connectivity (Wunsch, 2006), especially when many ‘stepping stones’ are involved (e.g. from Greenland ice via the North Atlantic Ocean to the European mainland).


Here, we test a different approach to compare D-O signals in Greenland ice cores (Andersen *et al.*, 2006) with abrupt events recorded in the high-resolution dated sediment sequence from the site Les Échets in southeastern France (Wohlfarth *et al.*, 2008). This multi-proxy lake record is dated with numerous radiocarbon and infrared stimulated luminescence (IRSL) measurements, and covers the time period ca. 45–15 ka BP. The record displays pronounced environmental changes which show striking similarities to D-O events in the Greenland ice cores. Instead of imposing synchronicity through tuning of the proxy events, we apply Bayesian techniques (Blaauw *et al.*, 2007a) to systematically compare the timing of the events recognised in the Les Échets sequence with those in the NGRIP ice core.

## Methods

Core EC1 was drilled in the deepest part of the former lake at Les Échets (LE;  $45^\circ 54' \text{ N}$ ,  $4^\circ 56' \text{ E}$ , 275 m above sea level) and analysed at high temporal resolution for multiple environmental proxies (Wohlfarth *et al.*, 2008). The alternating organic and inorganic sediments between 30.06 and 5 m depth were dated using 46 accelerator mass spectrometry (AMS) radiocarbon dates on plant remains, pollen grains and the insoluble sediment organic fraction, as well as 21 IRSL dates (Wohlfarth *et al.*, 2008). The major source of uncertainty for the  $^{14}\text{C}$  dates lies in the calibration to calendar ages, whereas for the IRSL dates uncertainty resides in the variability of the water content of the sediments. Differential compaction of the organic and inorganic sediments indicates that their water content changed during and after deposition. Therefore present-day moisture measurements were corrected using an estimated average water content of  $80 \pm 10\%$  over time.

Most of the LE radiocarbon ages are older than the limit of the IntCal04 calibration curve at 26 cal. ka BP (Reimer *et al.*, 2004). Although several curves for estimating calendar ages beyond that range are potentially available (Kitagawa and van der Plicht, 1998; Beck *et al.*, 2001; van der Plicht *et al.*, 2004; Fairbanks *et al.*, 2005; Hughen *et al.*, 2006; Danzeglocke *et al.*, 2008), we chose here the curves of Hughen *et al.* (2006, hereafter Hughen06) and Fairbanks *et al.* (2005, hereafter Fairbanks05), because of their long time-spans, high-dating resolution and reasonable between-curve agreements. Both Kitagawa and van der Plicht (1998) and Beck *et al.* (2001) provide alternative glacial curves with a reasonably high density, but both curves are in disagreement with each other and with Hughen06 and Fairbanks05 during at least parts of our period of interest. While the calendar ages of Hughen06 are based on tuning Cariaco Basin sediment palaeoclimate records to the high-resolution U/Th dated Hulu cave  $\delta^{18}\text{O}$  record (Wang *et al.*, 2001), those of Fairbanks05 were constructed from non-tuned, independently obtained U/Th dates on corals. Both curves have uncertainties in their calendar age models (errors 300–500 a for Hughen06, 50–100 a for Fairbanks05), and imprecisely known and possibly varying local marine reservoir effects (Muscheler *et al.*, 2008) create additional uncertainties in the radiocarbon ages of both curves. In order to take into account the uncertainties on the calendar as well as the radiocarbon age scales, we resampled each data point of the Hughen06 or Fairbanks05 curve assuming normally distributed errors of the radiocarbon and calendar age estimates, and repeated this process 10 000 times. For each iteration we interpolated linearly between the sampled  $^{14}\text{C}$  and cal. BP points, to produce  $^{14}\text{C}$  age point estimates at 50 calendar year intervals between 60 and 15 cal. ka BP. For each of these calendar years, histograms of all  $^{14}\text{C}$  age estimates were drawn to calculate 1 standard deviation (SD) highest posterior densities on the radiocarbon age scale. Calendar ages from the comparison curves produced above should not be considered as reliable as those derived from ‘official’ IntCal calibration curves, and thus we refrain from using the term ‘cal. BP’ (van der Plicht *et al.*, 2004).

The age models were constrained by prior information (for detailed explanations of the priors see Blaauw and Christen, 2005; Blaauw *et al.*, 2007b): (i) the depositional setting dictates that the dated depths must be ordered chronologically; (ii) given that 25 m of sediment had accumulated over ca. 25 ka, deposition rates of ca.  $5\text{--}20 \text{ cm}^{-1}$  are considered most likely, and other (positive) accumulation rates are possible although less likely (Fig. 1(a);  $\text{AlphaM} = 12$ ,  $\text{AlphaStd} = 4$ ); (iii) the variable stratigraphy suggests that rapid accumulation



**Figure 1** Distributions of the prior information for the Les Échets age–depth model. All are gamma distributions, which are comparable to the Gaussian distribution, although values are always positive and distributions can be asymmetric. The prior for the accumulation rate within a section is shown in (a), with a mean of 12 and a variation of 4 (AlphaM = 12, AlphaStd = 4). Between sections, accumulation rate can vary according to a gamma distribution centred around 1, with a variability set at epsilon = 2 (b). Hiatus sizes (c) are most likely to be very short (0 a or negligible), but stratigraphical information indicates hiatuses can occasionally last for several millennia (HiatusA =  $10^{-4}$ , HiatusB =  $10^{-5}$ )

rate changes and hiatuses could have occurred (Fig. 1(b); Epsilon = 2); (iv) between sections hiatuses are most likely to be short or absent, but could at times last millennia (Fig. 1(c); HiatusA =  $10^{-4}$ , HiatusB =  $10^{-5}$ ); and (v) every date has a prior probability of being outlying (5% for most (57) dates; 50% for seven  $^{14}\text{C}$  dates which could not be pre-treated, had a sample weight <1 mg or were possibly altered by a hard-water effect). Because there were no stratigraphical or other indications for a hiatus at 1653 and 1605 cm (as opposed to around 2700 cm; Wohlfarth *et al.*, 2008), we assumed that the two dates at those depths were erroneous, and adapted their prior outlier probabilities to 95%. The lowermost date (3097 cm depth) has not been pretreated and is therefore not considered to be reliable either. This part of the sediment sequence has several sand layers with sharp, erosive upper and lower boundaries, which indicate reworking and redeposition of sediments and likely also sedimentary hiatuses. This observation gives further evidence that the sediments in the lowest part of the analysed sequence might be older than suggested by the radiocarbon dates. Therefore we assigned 95% prior outlier probabilities (see below) to the lowermost two radiocarbon dates. Outlying dates were not removed but were identified and down-weighted automatically (see Blaauw and Christen, 2005). The IRSL and  $^{14}\text{C}$  dates of LE were combined with the above prior information, producing >1.7 billion Markov chain Monte Carlo age–depth models. Of these age model iterations, through thinning 10 000 were retained for subsequent age model analysis (Blaauw and Christen, 2005).

Most dates fit within 1 SD of the comparison curves (radiocarbon) or the 1:1 curve (IRSL), with the exceptions of (i) the lower part of the sequence where IRSL dates suggest much older ages than the radiocarbon dates, (ii) the upper part where IRSL dates suggest somewhat older ages than the radiocarbon dates, and (iii) outlying radiocarbon dates between ca. 37 and 26 ka BP. The Bayesian age-modelling process automatically identifies and down-weights outlying dates; mean posterior outlier probabilities were ~25% and 27% respectively for the Huguenot06 and Fairbanks05 runs. During the age modelling, all dates are used together while taking into account their stratigraphical ordering (Blaauw and Christen, 2005); as a result the age model confidence intervals are generally much narrower than those of the individual dates. Histograms of the calendar age distribution can be obtained for every depth through calculating the assigned calendar age for each of the 10 000 iterations, and graphed as grey-scales, where more likely calendar ages (appearing more frequently in the sampling process) are darker (Fig. 2(c)–(d)). The intervals


with conflicting IRSL and radiocarbon dates obtained much larger chronological uncertainties (wide grey areas). We therefore focus on the high-resolution dated middle part of the sequence, between 40 and 26 ka BP, where average 1 SD uncertainties are ca. 400 a.

The probability that an event took place in a proxy archive during a certain period depends on (i) the archive's chronological uncertainty (Fig. 2), (ii) the strength of the proxy evidence for an event (i.e. signal strength) (Fig. 3), and (iii) the duration of the period or window width (indicated by colours in Fig. 4). Here we assess all of these uncertainty sources together by calculating the event probability within a time window for the individual archives. The loss-on-ignition (LOI) and planktonic diatom proxies of LE were normalised to 100% and resampled assuming normally distributed errors (1% for LOI, 2% for diatoms), after which a running median was calculated (smoothing 25 for LOI, 11 for lower-resolution diatom data). All depths  $d$  where the running median increased beyond the error threshold were 'flagged'. This process was repeated 1000 times, after which for all depths of the individual proxies the ratios of flagged iterations were plotted (Fig. 3(a) and (b)). The means of the ratios of both proxies estimate the probabilities of enhanced lake productivity (Fig. 3(c)). Depths with probabilities <5% were neglected. For all age model iterations of LE or NGRIP, we find those depths  $d$ , with event probabilities  $p(e_d)$ , that fall within a time window with boundaries  $y_{\max}$  and  $y_{\min}$ . The probability that an event took place in an archive during this time window is equal to  $1 - \prod(1 - p(e_d))$ . In other words, we calculate the probability that *no* event took place within the time window, which is the product of  $1 - p(e_d)$ , and subtract this probability from 1 to find the probability that at least one event took place.

All calculations were performed in R (R Development Core Team, 2009; current version 2.9.0).

## Results

Intervals of higher lake organic productivity (LE events A–G in Fig. 3), interpreted to reflect warmer and more humid climatic conditions, are indicated by rising values for LOI and planktonic diatom concentrations, synchronously with many other analysed proxies in LE sediments (Ampel *et al.*, 2008; Wohlfarth *et al.*, 2008; Veres *et al.*, 2009). In contrast, low lake organic productivity, interpreted to reflect a colder and drier



**Figure 2** Bayesian age–depth models using two alternative radiocarbon comparison curves. The radiocarbon dates of Les Échets were matched against the Hughen06 (a, c) and Fairbanks05 (b, d) comparison curves. Posterior outlier probability is indicated by redness. Dashed curve is 1:1 line to which the IRSL dates (open diamonds) were matched. The age distributions are graphed as grey-scales (Blaauw *et al.*, 2007a) (c, d). Dark areas indicate secure sections of the age models, while lighter grey areas warn us of sections of a core where the chronological uncertainty is large. This figure is available in colour online at [www.interscience.wiley.com/journal/jqs](http://www.interscience.wiley.com/journal/jqs)


climate, is suggested by low values of LOI and planktonic diatom concentrations. Most of the LE warm events were accompanied by clear increases in arboreal pollen within the same core (Wohlfarth *et al.*, 2008), suggesting that the environmental changes were regional at least. Peaks A–D clearly denote events of increased lake organic productivity (high event probabilities), whereas events E–G appear more subtle. Wohlfarth *et al.* (2008) suggested that the shifts between higher and low lake organic productivity phases resemble D–O and Greenland Interstadial (GI) events in the Greenland ice cores. Although it would be tempting to tune these intervals by eye to GI events 8–3 (Fig. 3), instead we calculate the statistical probability that both independently dated archives reacted simultaneously within certain time windows, taking into account their chronological uncertainties (Blaauw *et al.*, 2007a).

Significant LE event probabilities (95%) are only reached by employing time windows of 400 a or wider. The choice of  $^{14}\text{C}$  comparison curve has a considerable effect on the timing of events A–D in particular, owing to disagreements between Hughen06 and Fairbanks05 around 37–34 ka BP. This disagreement is partly caused by Fairbanks05 having fewer evenly spaced data points for this time interval, and thus more kinks than Hughen06. Overlapping, closely spaced GI events (Fig. 4(c)) can hardly be distinguished from each other on an absolute timescale. Although the relative timing between GI events is well known (see below), NGRIP cumulative chronological uncertainties (Andersen *et al.*, 2006) prevent us from assigning an individual GI event to any particular decade or even century (>95% event probabilities using time windows of 1550 a or wider).

To test for synchronous events between LE and NGRIP within a time window, we can simply calculate the product of the

event probabilities from the individual archives (Fig. 4(d) and (e)). High probabilities indicate that an event of increased lake organic productivity in Les Échets likely occurred simultaneously with a GI event, whereas low probabilities indicate the absence of a simultaneous reaction and/or a lack of information (e.g. hiatus, large chronological uncertainties). Probabilities of synchronous events between LE and NGRIP never reach 100% at centennial resolutions. For example, only within >1000 a time windows does it seem likely (95%) that event LE–G co-occurred with one of the closely spaced GI events 4 or 3 in NGRIP. The other events show significant synchronicity using time windows of 1250 a or wider. In all cases, synchronous reactions at decadal to multi-centennial year resolution are very unlikely, since probabilities never exceed 10% (red shades in Fig. 4(d) and (e)). Wider window sizes tend to reach higher event probabilities because (i) the windows include more of the probability distributions of individual events, and (ii) probability distributions of multiple separate events could cover the time windows considered.

Since the NGRIP timescale is based on annual layer counting, *absolute* chronological uncertainties accumulate to millennial scales over time during the glacial. However, cumulative layer-count errors between neighbouring GI events within NGRIP are much smaller, reaching centennial to decadal scales (Andersen *et al.*, 2006). Therefore, the *relative* timings between GI events 8 to 3 are much more tightly constrained than those between the IRSL/ $^{14}\text{C}$  dated events LE–A to LE–G. Given the large absolute chronological uncertainties on the timing of individual events (Fig. 4), we investigate the patterns of timing between events (Fig. 5). Even if the absolute timings of events in either NGRIP or LE are erroneous, with a common climate mechanism forcing both the NGRIP and LE events, a shared ‘fingerprint’ of the timing between events



**Figure 3** Les Échets proxies plotted against depth (black lines) and compared with the NGRIP  $\delta^{18}\text{O}$  record (Dansgaard *et al.*, 1993). Probabilities of enhanced lake productivity in Les Échets were calculated, through finding those depths with major increases in LOI (a) or planktonic diatoms (b) (see text). Clear increases of the individual proxies are plotted as grey histograms. The means of these proxy increases estimate the probabilities of enhanced lake productivity (histograms in c). The events A–G of enhanced lake productivity could be tuned (grey connecting lines show possible links) to abrupt  $\delta^{18}\text{O}$  rises in NGRIP (Andersen *et al.*, 2006), Greenland Isotope (GI) events 8 to 3 (d). Horizontal bars in (d) indicate 1 SD dating confidence intervals of the GI events. This figure is available in colour online at [www.interscience.wiley.com/journal/jqs](http://www.interscience.wiley.com/journal/jqs)

should be expected (but for similar event frequencies in unrelated climate series see Wunsch, 2006). Whereas the timing between LE events F and G is comparable to that between GI events 4 and 3, the relative timings between the other events differ by many centuries. Moreover, there appear to be more LE than GI events in the studied period.

We could postulate several reasons for the offsets between the timing of events between Les Échets and NGRIP, e.g. erroneous dates, large changes in marine reservoir age for the Huguen06 and Fairbanks05 curves, unrecognised hiatuses, low resolution, erosion and/or reworking of older layers in the LE record, missing or false proxy responses to climate change in either archive, or hundreds of 'false' years in the NGRIP timescale (e.g. see Skinner, 2008). However, our approach shows that the assumption of synchronous events between both archives might be erroneous; the alternative explanation that the events were asynchronous should not be discarded.

## Discussion


Our results appear to deviate from an apparently sound story based on a large body of stratigraphical studies (Bond *et al.*, 1993; Cacho *et al.*, 1999; Shackleton *et al.*, 2000; Sánchez-Goñi *et al.*, 2002; Rohling *et al.*, 2003; Tzedakis *et al.*, 2004; Roucoux *et al.*, 2005; Skinner, 2008). It could be argued that our failure to find simultaneous events between Greenland and

a terrestrial site in Europe was caused by proxy or dating problems (see above), and thus should not lead us to call into question the reported tele-connections between Greenland and other proxy archives. For example, within Mediterranean marine cores located relatively near to Les Échets, sea surface temperature and arboreal pollen content changed in concert (Sánchez-Goñi *et al.*, 2002). Interestingly enough, the arboreal pollen records also show a close visual resemblance to that of Les Échets (Wohlfarth *et al.*, 2008), indicating spatially widespread events of climate change. As the events in the Mediterranean cores were reportedly simultaneous with events in Greenland (Sánchez-Goñi *et al.*, 2002), it should follow that Les Échets must have reacted simultaneously as well. However, the synchronicity of the Mediterranean climate events to those in Greenland is not surprising, since their chronologies were based on aligning foraminifera  $\delta^{18}\text{O}$  and sea surface temperature fluctuations with  $\delta^{18}\text{O}$  peaks in the GISP2 ice core record.

Asynchronous climate events between proxy archives do not necessarily require us to invoke leads or lags, although these can be calculated and tested easily using our time window method. Indeed, probably the safest (though most pessimistic) 'null hypothesis' for any proxy-derived climate event would be to assume that it is unrelated to other events, until proven otherwise by independent, replicated data. However, millennial-scale chronological uncertainties for most of the last glacial period have prevented us from directly proving synchronicity of D–O events through independently dated archives. Over the years a plausible history of D–O events has been reconstructed, based on climatic reasoning combined with event ages derived from tuning between archives. It should not be forgotten though that some degree of circular reasoning has necessarily been involved in this process. Indeed, it has been known for decades that assuming synchronicity of past climate/environmental events without independent confirmation (Walker *et al.*, 1999) can lead to a 'reinforcement syndrome' (von Post, 1946; Smith and Pilcher, 1973; Baillie, 1991; Oldfield, 2001; Bennett, 2002).

Past climate research would benefit from a better quantification of the uncertainties involved. Until now, most tuning has been performed visually, involving subjective decisions. Statistical methods could be developed to better quantify the uncertainties of tuned age models, e.g. based on (multi-proxy, multi-site) shape comparisons of events. In the longer run, one could envisage obtaining a better understanding of D–O events through a coupled ice/ocean/atmosphere/biosphere climate model guided by a spatial array of independently dated proxy events, while taking into account their chronological and other uncertainties. A step in this direction can be made by applying our time window approach to a larger set of independently dated marine and/or terrestrial proxy archives.


The abrupt temperature fluctuations reconstructed from Greenland ice cores (Dansgaard *et al.*, 1993; Steffensen *et al.*, 2008) must have been forced by major climatic events, which arguably also should have had effects on larger spatial scales (Wunsch, 2006). However, most of the reported synchronous climate events outside Greenland were tuned to GI events (Bond *et al.*, 1993; Cacho *et al.*, 1999; Shackleton *et al.*, 2000; Sánchez-Goñi *et al.*, 2002; Tzedakis *et al.*, 2004; Roucoux *et al.*, 2005; Skinner, 2008) and thus cannot themselves be used to infer synchronicity. Even with the highest-resolution dated age models (67 radiocarbon and IRSL dates for Les Échets, multi-proxy annual layer counting for Greenland; Andersen *et al.*, 2006), chronological uncertainties are currently too high to resolve through independent chronologies whether last glacial D–O climate events were simultaneous, or even related, between Greenland and western Europe. These problems are even more serious with proxy archives dated and analysed at



**Figure 4** Events of enhanced lake productivity in Les Échets from age–depth models using either Hughen06 (a) or Fairbanks05 (b), compared with GI events in NGRIP (c). Probabilities of events of enhanced lake productivity (Les Échets) or temperature rises in Greenland (GI events from Andersen *et al.*, 2006) were calculated using a time window approach modified from Blaauw *et al.* (2007a) (see text). Time windows moved at 50 a jumps. Window sizes were 50–2000 a in steps of 50 a, shown as rainbow colours in the probability histograms (colour legend for window sizes is shown at bottom left). The histograms are centred at the mid-points of the time windows, e.g. the probability of an event between 40 and 38 ka BP is shown as the height of the violet colour at 39 ka BP. Histograms of narrower window sizes are plotted overlaying those of wider windows sizes (narrower histograms reach lower probabilities). The probabilities of simultaneous events between Les Échets and NGRIP within time windows are shown in (d, e)

lower resolution. Although one would wish to resolve the spatiotemporal nature of past glacial climate events at annual or, at most, decadal scales, currently we are limited to comparisons at multi-centennial to millennial resolution only. However, this problem could be reduced significantly if more widespread time markers such as well-defined tephra layers were to be found.

**Acknowledgements** BW acknowledges support from the Swedish Research Council (VR). MB thanks Keith Bennett for help at various stages of this paper. Comments from William D’Andrea, Maria Sánchez Goñi, John Lowe, Chronis Tzedakis and an anonymous referee on a previous version of this paper have helped improve it considerably. This is a contribution to ESF EuroCores on EuroCLIMATE project RESOLuTION.



**Figure 5** Timing in calendar years between neighbouring events of the Les Échets and NGRIP records. Colour shading shows the timing between Les Échets A–G using Hughen05 (green) or Fairbanks05 (red), and NGRIP GI events 8 to 3 (blue). Darker colours indicate more likely timing. Uncertainties of NGRIP timing are based on the accumulated counting error between neighbouring events (Andersen *et al.*, 2006). There seems to be one more event in LE than in NGRIP during the period of study, which causes additional confusion when deciding which events to correlate

## References

- Allen JRM, Brandt U, Brauer A, Hubberten H-W, Huntley B, Keller J, Kram M, Mackensen A, Mingram J, Negendank JFW, Nowaczyk NR, Oberhänsli H, Watts WA, Wulf S, Zolitschka B. 1999. Rapid environmental changes in southern Europe during the last glacial period. *Nature* **400**: 740–743.
- Ampel L, Wohlfarth B, Risberg J, Veres D. 2008. Paleolimnological response to millennial and centennial scale climate variability during MIS 3 and 2 as suggested by the diatom record in Les Échets, France. *Quaternary Science Reviews* **27**: 1493–1504.
- Andersen KK, Svensson A, Johnsen SJ, Rasmussen SO, Bigler M, Röhlisberger R, Ruth U, Siggaard-Andersen ML, Steffensen JP, Dahl-Jensen D, Vinther BM, Clausen HB. 2006. The Greenland ice core chronology 2005, 15–42 ka. Part 1: Constructing the time scale. *Quaternary Science Reviews* **25**: 3246–3257.
- Austin WEN, Wilson LJ, Hunt JB. 2004. The age and chronostratigraphical significance of North Atlantic Ash Zone II. *Journal of Quaternary Science* **19**: 137–146.
- Baillie MGL. 1991. Suck in and smear: two related chronological problems for the 90s. *Journal of Theoretical Archaeology* **2**: 12–16.
- Beck JW, Richards DA, Edwards RL, Silverman BW, Smart PL, Donahue DJ, Hererra-Osterheld S, Burr GS, Calsoyas L, Jull AJT, Biddulph D.

2001. Extremely large variations of atmospheric C-14 concentration during the last glacial period. *Science* **292**: 2453–2458.
- Bennett KD. 2002. Comment: the Greenland 8200 cal. yr BP event detected in loss-on-ignition profiles in Norwegian lacustrine sediment sequences. *Journal of Quaternary Science* **17**: 97–99.
- Blaauw M, Christen JA. 2005. Radiocarbon peat chronologies and environmental change. *Applied Statistics* **54**: 805–816.
- Blaauw M, Christen JA, Mauquoy D, van der Plicht J, Bennett KD. 2007a. Testing the timing of radiocarbon-dated events between proxy archives. *The Holocene* **17**: 283–288.
- Blaauw M, Bakker R, Christen JA, Hall VA, van der Plicht JA. 2007b. Bayesian framework for age-modelling of radiocarbon dated peat deposits: case studies from the Netherlands. *Radiocarbon* **49**: 357–367.
- Bond G, Broecker W, Johnsen S, McManus J, Labeyrie L, Jouzel J, Bonani G. 1993. Correlations between climate records from North Atlantic sediments and Greenland ice. *Nature* **365**: 143–147.
- Burns SJ, Fleitmann D, Matter A, Al-Subbaray AA. 2003. Indian Ocean climate and an absolute chronology over Dansgaard/Oeschger events 9 to 13. *Science* **301**: 1365–1367.
- Cacho I, Grimalt JO, Pelejero C, Canals M, Sierro FJ, Flores JA, Shackleton N. 1999. Dansgaard–Oeschger and Heinrich event imprints in Alboran Sea paleotemperatures. *Paleoceanography* **14**: 698–705.
- Dansgaard W, Johnsen SJ, Clausen HB, Dahl-Jensen D, Gundestrup NS, Hammer CU, Hvidberg CS, Steffensen JP, Sveinbjörnsdóttir AE, Jouzel J, Bond G. 1993. Evidence for general instability of past climate from a 250-kyr ice-core record. *Nature* **364**: 218–220.
- Danzeglocke U, Jöris O, Weninger B. 2008. CalPal-2007<sup>online</sup>. <http://www.calpal-online.de> [23 March 2009].
- EPICA Community Members. 2006. One-to-one coupling of glacial climate variability in Greenland and Antarctica. *Nature* **444**: 195–198.
- Fairbanks RG, Mortlock RA, Chiu TC, Cao T, Kaplan A, Guilderson TP, Fairbanks TW, Bloom AL, Grootes PM, Nadeau MJ. 2005. Radiocarbon calibration curve spanning 0 to 50,000 years BP based on paired <sup>230</sup>Th/<sup>234</sup>U/<sup>238</sup>U and <sup>14</sup>C dates on pristine corals. *Quaternary Science Reviews* **24**: 1781–1796.
- Hendy IL, Kennett JP, Roark EB, Ingram BL. 2002. Apparent synchrony of submillennial scale climate events between Greenland and Santa Barbara Basin, California from 30–10 ka. *Quaternary Science Reviews* **21**: 1167–1184.
- Hughen K, Lehman S, Southon J, Overpeck J, Marshal O, Herring C, Turnbull J. 2004. <sup>14</sup>C activity and global carbon cycle changes over the past 50,000 years. *Science* **303**: 202–207.
- Hughen K, Southon J, Lehman S, Bertrand C, Turnbull J. 2006. Marine-derived <sup>14</sup>C calibration and activity record for the past 50,000 years updated from the Cariaco Basin. *Quaternary Science Reviews* **25**: 3216–3227.
- Kitagawa H, van der Plicht J. 1998. Atmospheric radiocarbon calibration to 45,000 yr B.P.: late glacial fluctuations and cosmogenic isotope production. *Science* **279**: 1187–1190.
- Knutti R, Flückiger J, Stocker TF, Timmermann A. 2004. Strong hemispheric coupling of glacial climate through freshwater discharge and ocean circulation. *Nature* **430**: 851–856.
- Liesiecki LE, Raymo ME. 2005. A Pliocene–Pleistocene stack of 57 globally distributed benthic  $\delta^{18}\text{O}$  records. *Paleoceanography* **20**: PA1003.
- Martinson DG, Pisias NG, Hays JD, Imbrie J, Moore TC Jr, Shackleton NJ. 1987. Age dating and the orbital theory of the ice ages: development of a high-resolution 0 to 300,000-year chronostratigraphy. *Quaternary Research* **27**: 1–29.
- Muscheler R, Kromer B, Björck S, Svensson A, Friedrich M, Kaiser KF, Southon J. 2008. Tree rings and ice cores reveal <sup>14</sup>C calibration uncertainties during the Younger Dryas. *Nature Geoscience* **1**: 263–267.
- Oldfield F. 2001. A question of timing: a comment on Hong, Jiang, Liu, Zhou, Beer, Li, Leng, Hong and Qin. *The Holocene* **11**: 123–124.
- R Development Core Team. 2009. *R: a language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria (<http://www.R-project.org>).
- Reimer PJ, Baillie MGL, Bard E, Bayliss A, Beck JW, Bertrand CJH, Blackwell PG, Buck CE, Burr GS, Cutler KB, Damon PE, Edwards RL, Fairbanks RG, Friedrich M, Guilderson TP, Hogg AG, Hughen KA, Kromer B, McCormac G, Manning S, Bronk Ramsey C, Remmele RWRS, Southon JR, Stuiver M, Talamo S, Taylor FW, van der Plicht J, Weyhenmeyer CE. 2004. INTCAL04 terrestrial radiocarbon age calibration, 0–26 cal kyr BP. *Radiocarbon* **46**: 1029–1058.
- Rohling EJ, Mayewski PA, Challenor P. 2003. On the timing and mechanism of millennial-scale climate variability during the last glacial cycle. *Climate Dynamics* **20**: 257–267.
- Roucoux KH, de Abreu L, Shackleton NJ, Tzedakis PC. 2005. The response of NW Iberian vegetation to North Atlantic climate oscillations during the last 65 kyr. *Quaternary Science Reviews* **24**: 1637–1653.
- Sánchez-Goni MF, Cacho I, Turon JL, Guiot J, Sierro FJ, Peypougoet JP, Grimalt JO, Shackleton NJ. 2002. Synchronicity between marine and terrestrial responses to millennial scale climatic variability during the last glacial period in the Mediterranean region. *Climate Dynamics* **19**: 95–105.
- Shackleton NJ, Hall MA, Vincent E. 2000. Phase relationships between millennial-scale events 64,000–24,000 years ago. *Paleoceanography* **15**: 565–569.
- Shackleton NJ, Fairbanks RG, Chiu T-C, Parrenin F. 2004. Absolute calibration of the Greenland time scale: implications for Antarctic time scales and for D<sup>14</sup>C. *Quaternary Science Reviews* **23**: 1513–1522.
- Skinner LC. 2008. Revisiting the absolute calibration of the Greenland ice-core age-scales. *Climate of the Past* **4**: 295–302.
- Smith AG, Pilcher JR. 1973. Radiocarbon dates and vegetational history of the British Isles. *New Phytologist* **72**: 903–914.
- Steffensen JP, Andersen KK, Bigler M, Clausen HB, Dahl-Jensen D, Fischer H, Goto-Azuma K, Hansson M, Johnsen SJ, Jouzel J, Masson-Delmotte V, Popp T, Rasmussen SO, Röthlisberger R, Ruth U, Stauffer B, Siggaard-Andersen M-L, Sveinbjörnsdóttir ÁE, Svensson A, White JWC. 2008. High-Resolution Greenland ice core data show abrupt climate change happens in few years. *Science* **321**: 680–684.
- Svensson A, Andersen KK, Bigler M, Clausen HB, Dahl-Jensen D, Davies SM, Johnsen SJ, Muscheler R, Rasmussen SO, Röthlisberger R, Steffensen JP, Vinther BM. 2006. The Greenland ice core chronology 2005, 15–42 ka. Part 2: Comparison to other records. *Quaternary Science Reviews* **25**: 3258–3267.
- Svensson A, Andersen KK, Bigler M, Clausen HB, Dahl-Jensen D, Davies SM, Johnsen SJ, Muscheler R, Parrenin F, Rasmussen SO, Röthlisberger R, Seierstad I, Steffensen JP, Vinther BM. 2008. A 60 000 year Greenland stratigraphic ice core chronology. *Climate of the Past* **4**: 47–57.
- Turney CSM, Kershaw AP, Clemens SC, Branch N, Moss PT, Fifield LK. 2004. Millennial and orbital variations of El Niño/Southern Oscillation and high-latitude climate in the last glacial period. *Nature* **428**: 306–310.
- Tzedakis PC, Frogley MR, Lawson IT, Preece RC, Cacho RC, de Abreu L. 2004. Ecological thresholds and patterns of millennial-scale climate variability: the response of vegetation in Greece during the last glacial period. *Geology* **32**: 109–112.
- van der Plicht J, Beck JW, Bard E, Baillie MGL, Blackwell PG, Buck CE, Friedrich M, Guilderson TP, Hughen KA, Kromer B, McCormac G, Ramsey CB, Reimer PJ, Reimer RW, Remmele S, Richards DA, Southon JR, Stuiver M, Weyhenmeyer CE. 2004. NotCal04: comparison/calibration <sup>14</sup>C records 26–50 cal. kyr BP. *Radiocarbon* **46**: 1225–1238.
- Vautravers MJ, Shackleton NJ. 2006. Centennial-scale surface hydrology off Portugal during marine isotope stage 3: insights from planktonic foraminiferal fauna variability. *Paleoceanography* **21**: PA3004.
- Veres D, Lallier-Vergès E, Wohlfarth B, Lacourse T, Kéravis D, Björck S, Preusser F, Andrieu-Ponel V, Ampel L. 2009. Climate-driven changes in lake conditions during late MIS 3 and MIS 2: a high-resolution geochemical record from Les Échets, France. *Boreas* **38**: 230–243.
- Voelker AHL and workshop participants. 2002. Global distribution of centennial-scale records for Marine Isotope Stage (MIS) 3: a data base. *Quaternary Science Reviews* **21**: 1185–1212.
- Von Post L. 1946. The prospect for pollen analysis in the study of the Earth's climatic history. *New Phytologist* **45**: 198–203.
- Walker MJC, Björck S, Lowe JJ, Cwynar LC, Johnsen S, Knudsen K-L, Wohlfarth B. INTIMATE Group. 1999. Isotopic ‘events’ in the GRIP

- ice core: a stratotype for the late Pleistocene. *Quaternary Science Reviews* **18**: 1143–1150.
- Wang Y, Cheng H, Edwards RL, An Z, Wu J, Shen CC, Dorale JA. 2001. A high-resolution absolute-dated monsoon record from Hulu Cave, China. *Science* **294**: 2345–2348.
- Wåstegård S, Rasmussen TL, Kuijpers A, Nielsen T, van Weering TCE. 2006. Composition and origin of ash zones from Marine Isotope Stages 3 and 2 in the North Atlantic. *Quaternary Science Reviews* **25**: 2409–2419.
- Wohlfarth B, Veres D, Ampel L, Lacourse T, Blaauw M, Preusser F, Andrieu-Ponel V, Kéravis D, Lallier-Vergès E, Björck S, Davies SM, de Beaulieu J-L, Risberg J, Hormes A, Kasper HU, Possnert G, Reille M, Thouveny N, Zander A. 2008. Rapid ecosystem response to abrupt climate changes during the last glacial period in western Europe, 40–16 ka. *Geology* **36**: 407–410.
- Wunsch C. 2006. Abrupt climate change: an alternative view. *Quaternary Research* **65**: 191–203.